



ISAKOS
CONGRESS
2025



MUNICH
GERMANY
June 8-11

Does scaffold enhancement show significant superiority over microfracture alone for treating knee chondral defects? A systematic review and meta-analysis of randomised clinical trials.

Marta Kubisa, Karol Pałka, Anna Akbas, Michał Kubisa, Michał Dobrakowski



Faculty Disclosure Information

- Nothing to disclosure
- Carolina Hospital - Warsaw, Poland



ISAKOS
CONGRESS
2025



MUNICH
GERMANY
June 8-11

PURPOSE

Chondral and osteochondral lesions in the knee are common conditions that significantly impair individuals' well-being and can lead to osteoarthritis, imposing substantial burdens on healthcare systems. The limited natural healing capacity of articular cartilage necessitates innovative treatment strategies. Microfracture (MF) is a widely used technique for knee chondral defects, but its long-term efficacy is often inadequate. Although recent randomised controlled trials have compared microfractures with scaffold-enhanced therapies, a comprehensive systematic review and meta-analysis are lacking.




ISAKOS
CONGRESS
2025



MUNICH
GERMANY
June 8–11

METHODS

An extensive literature search was conducted in PubMed and EMBASE databases following PRISMA guidelines. Inclusion criteria focused on randomised controlled trials (RCTs) comparing microfractures alone to matrix-induced chondrogenesis for knee chondral defects with at least a 12-month follow-up. Ten randomised controlled trials conducted between 2013 and 2024, enrolling 378 patients, were included.



RESULTS

Cohort studies included in meta-analysis and the results of the modified Coleman Methodology Score (mCMS)

Authors	Anders et al. [2]	Chung et al. [10]	Kon et al. [30]	Volz et al. [51]	Kim et al. [27]	Kim et al. [26]	Shive et al. [43]	Stanish et al. [44]	Glasbrenner et al. [18]	Altschuler et al. [1]
Part A										
1. Study size	7	10	10	7	4	10	10	10	4	10
2. Mean duration of follow-up	2	2	2	5	2	2	5	2	5	2
3. Number of treatment procedures	10	10	10	10	10	10	10	10	10	10
4. Type of study	15	15	15	15	15	15	15	15	15	15
5. Diagnostic certainty	5	5	5	5	5	5	5	5	5	5
6. Description of surgical procedure	5	5	5	5	5	5	5	5	5	5
7. Description of rehabilitation procedures	10	10	5	10	5	5	10	10	10	10
Part B										
1. Outcome criteria										
1a	2	2	2	2	2	2	2	2	2	2
1b	2	2	2	2	2	2	2	2	2	2
1c	4	4	4	4	4	4	4	4	4	4
2. Procedure for assessing outcomes										
2a	5	5	5	5	5	5	5	5	5	5
2b		4	4		4	4		4		
2c	3	3	3	3	3	3	3	3	3	3
2d	3	3	3	3	3	3	3	3	3	3
3. Subject selection process										
3a	5	5	5	5	5	5	5	5	5	
3b										
3c										
Total Part A (max 60)	54	57	52	57	46	52	60	57	54	57
Total Part B (max 40)	24	28	28	24	28	28	24	28	24	19
Total mCMS	78	85	80	81	74	80	84	85	78	76




ISAKOS
CONGRESS
2025





MUNICH
GERMANY
June 8–11

The meta-analysis showed no significant superiority of scaffoldover MF ($p > 0.05$) in International Knee Documentation Committee, Knee Injury and Osteoarthritis Outcome, Visual Analog Scale, and Magnetic Resonance Observation of Cartilage Repair Tissue scores at 12 and 24 months. However, individual studies suggested the potential benefits of scaffolds, especially in long-term outcomes. Clinical improvements from MF typically decline after 2–3 years, underscoring the need for long-term follow-up in future research.



MOCART -magnetic resonance observation of cartilage repair tissue





IKDC - International Knee Documentation Committee Scores



KOOS - Knee Injury and Osteoarthritis Outcome Scores



VAS - Visual Analog Scale



CONCLUSION

Our meta-analysis shows no significant difference between MF and MF with scaffold in treating knee cartilage defects, though some long-term RCTs demonstrate statistically significant differences. The absence of a universally accepted algorithm for analysing knee chondral defects limits this study. Establishing reliable guidelines and standardised study protocols is essential to improve long-term patient outcomes and the quality of future papers.



ISAKOS
CONGRESS
2025



MUNICH
GERMANY
June 8–11

1. Heir S, Nerhus TK, Løttersud JH, Løken S, Ekeland A, Engebretsen L, i in. Focal Cartilage Defects in the Knee Impair Quality of Life as Much as Severe Osteoarthritis: A Comparison of Knee Injury and Osteoarthritis Outcome Score in 4 Patient Categories Scheduled for Knee Surgery. Am J Sports Med. luty 2010;38(2):231–7.

2. Hjelle K, Solheim E, Strand T, Muri R, Brittberg M. Articular cartilage defects in 1,000 knee arthroscopies. Arthrosc J Arthrosc Relat Surg. wrzesień 2002;18(7):730–4.

3. Ding C, Cicuttini F, Scott F, Cooley H, Boon C, Jones G. Natural History of Knee Cartilage Defects and Factors Affecting Change. Arch Intern Med. 27 marzec 2006;166(6):651.

4. Gratz KR, Wong BL, Bae WC, Sah RL. The effects of focal articular defects on cartilage contact mechanics. J Orthop Res. maj 2009;27(5):584–92.

5. Heijink A, Gomoll AH, Madry H, Drobníč M, Filardo G, Espregueira-Mendes J, i in. Biomechanical considerations in the pathogenesis of osteoarthritis of the knee. Knee Surg Sports Traumatol Arthrosc. marzec 2012;20(3):423–35.

6. Steinwachs MR, Guggi Th, Kreuz PC. Marrow stimulation techniques. Injury. kwiecień 2008;39(1):26–31.

7. Marcacci M, Filardo G, Kon E. Treatment of cartilage lesions: What works and why? Injury. styczeń 2013;44:S11–5.

8. Goyal D, Keyhani S, Lee EH, Hui JHP. Evidence-Based Status of Microfracture Technique: A Systematic Review of Level I and II Studies. Arthrosc J Arthrosc Relat Surg. wrzesień 2013;29(9):1579–88.

9. Matsushita T, Tokura T, Okimura K, Sano S, Nishida K, Nagai K, i in. Surgical treatment of cartilage lesions in the knee: A narrative review. J Jt Surg Res. grudzień 2023;1(1):70–9.

10. Behrens P, Bitter T, Kurz B, Russlies M. Matrix-associated autologous chondrocyte transplantation/implantation (MACT/MACI)—5-year follow-up. The Knee. czerwiec 2006;13(3):194–202.

11. Brittberg M. Autologous Chondrocyte Transplantation. Clin Orthop. (3678).

12. Bentley G, Biant LC, Carrington RWJ, Akmal M, Goldberg A, Williams AM, i in. A prospective, randomised comparison of autologous chondrocyte implantation versus mosaicplasty for osteochondral defects in the knee. J Bone Joint Surg Br. marzec 2003;85-B(2):223–30.

13. Mithoefer K, McAdams T, Williams RJ, Kreuz PC, Mandelbaum BR. Clinical Efficacy of the Microfracture Technique for Articular Cartilage Repair in the Knee: An Evidence-Based Systematic Analysis. Am J Sports Med. październik 2009;37(10):2053–63.

14. Kim MS, Koh IJ, Choi YJ, Pak KH, In Y. Collagen Augmentation Improves the Quality of Cartilage Repair After Microfracture in Patients Undergoing High Tibial Osteotomy: A Randomized Controlled Trial. Am J Sports Med. lipiec 2017;45(8):1845–55.

15. Anders S, Volz M, Frick H, Gellissen J. A Randomized, Controlled Trial Comparing Autologous Matrix-Induced Chondrogenesis (AMIC®) to Microfracture: Analysis of 1- and 2-Year Follow-Up Data of 2 Centers. Open Orthop J. 3 maj 2013;7(1):133–43.

16. Volz M, Schaumburger J, Frick H, Grifka J, Anders S. A randomized controlled trial demonstrating sustained benefit of Autologous Matrix-Induced Chondrogenesis over microfracture at five years. Int Orthop. kwiecień 2017;41(4):797–804.

17. Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, i in. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. BMJ. 29 marzec 2021;n71.

18. Coleman BD, Khan KM, Maffulli N, Cook JL, Wark JD. Studies of surgical outcome after patellar tendinopathy: clinical significance of methodological deficiencies and guidelines for future studies. Scand J Med Sci Sports. luty 2000;10(1):2–11.

19. Ramponi L, Yasui Y, Murawski CD, Ferkel RD, DiGiovanni CW, Kerkhoffs GMMJ, i in. Lesion Size Is a Predictor of Clinical Outcomes After Bone Marrow Stimulation for Osteochondral Lesions of the Talus: A Systematic Review. Am J Sports Med. czerwiec 2017;45(7):1698–705.

20. Chung JY, Lee D hyung, Kim TH, Kwack KS, Yoon KH, Min BH. Cartilage extra-cellular matrix biomembrane for the enhancement of microfractured defects. Knee Surg Sports Traumatol Arthrosc. czerwiec 2014;22(6):1249–59.

21. Kon E, Filardo G, Brittberg M, Busacca M, Condello V, Engebretsen L, i in. A multilayer biomaterial for osteochondral regeneration shows superiority vs microfractures for the treatment of osteochondral lesions in a multicentre randomized trial at 2 years. Knee Surg Sports Traumatol Arthrosc. wrzesień 2018;26(9):2704–15.

22. Kim JH, Heo JW, Lee DH. Clinical and Radiological Outcomes After Autologous Matrix-Induced Chondrogenesis Versus Microfracture of the Knee: A Systematic Review and Meta-analysis With a Minimum 2-Year Follow-up. Orthop J Sports Med. 1 listopad 2020;8(11):232596712095928.

23. Shive MS, Stanish WD, McCormack R, Forriol F, Mohtadi N, Pelet S, i in. BST-CarGel® Treatment Maintains Cartilage Repair Superiority over Microfracture at 5 Years in a Multicenter Randomized Controlled Trial. CARTILAGE. kwiecień 2015;6(2):62–72.

24. Stanish WD, McCormack R, Forriol F, Mohtadi N, Pelet S, Desnoyers J, i in. Novel Scaffold-Based BST-CarGel Treatment Results in Superior Cartilage Repair Compared with Microfracture in a Randomized Controlled Trial. J Bone Jt Surg. 18 wrzesień 2013;95(18):1640–50.

25. Glasbrenner J, Petersen W, Raschke MJ, Steiger M, Verdonk R, Castelli CC, i in. Matrix-Augmented Bone Marrow Stimulation With a Polyglycolic Acid Membrane With Hyaluronan vs Microfracture in Local Cartilage Defects of the Femoral Condyles: A Multicenter Randomized Controlled Trial. Orthop J Sports Med. 1 maj 2020;8(5):232596712092293.

26. Altschuler N, Zaslav KR, Di Matteo B, Sherman SL, Gomoll AH, Hacker SA, i in. Aragonite-Based Scaffold Versus Microfracture and Debridement for the Treatment of Knee Chondral and Osteochondral Lesions: Results of a Multicenter Randomized Controlled Trial. Am J Sports Med. marzec 2023;51(4):957–67.

27. Knutsen G, Drogset JO, Engebretsen L, Grøntvedt T, Ludvigsen TC, Løken S, i in. A Randomized Multicenter Trial Comparing Autologous Chondrocyte Implantation with Microfracture: Long-Term Follow-up at 14 to 15 Years. J Bone Jt Surg. 17 sierpień 2016;98(16):1332–9.

28. Kraeutler MJ, Belk JW, Purcell JM, McCarty EC. Microfracture Versus Autologous Chondrocyte Implantation for Articular Cartilage Lesions in the Knee: A Systematic Review of 5-Year Outcomes. Am J Sports Med. marzec 2018;46(4):995–9.

29. Gobbi A, Karnatzikos G, Kumar A. Long-term results after microfracture treatment for full-thickness knee chondral lesions in athletes. Knee Surg Sports Traumatol Arthrosc. wrzesień 2014;22(9):1986–96.

30. Nehrer S, Spector M, Minas T. Histologic Analysis of Tissue After Failed Cartilage Repair Procedures: Clin Orthop. sierpień 1999;365:149–62.

31. Mithoefer K, Venugopal V, Manaqibwala M. Incidence, Degree, and Clinical Effect of Subchondral Bone Overgrowth After Microfracture in the Knee. Am J Sports Med. sierpień 2016;44(8):2057–63.

32. Tampieri A, Sandri M, Landi E, Pressato D, Francioli S, Quarto R, i in. Design of graded biomimetic osteochondral composite scaffolds. Biomaterials. wrzesień 2008;29(26):3539–46.

33. Serre CM, Papillard M, Chavassieux P, Voegel JC, Boivin G. Influence of magnesium substitution on a collagen-apatite biomaterial on the production of a calcifying matrix by human osteoblasts. J Biomed Mater Res. 15 grudzień 1998;42(4):626–33.

34. Kon E, Delcogliano M, Filardo G, Fini M, Giavaresi G, Francioli S, i in. Orderly osteochondral regeneration in a sheep model using a novel nano-composite multilayered biomaterial. J Orthop Res. styczeń 2010;28(1):116–24.

35. Delcogliano M, De Caro F, Scaravella E, Ziveri G, De Biase CF, Marotta D, i in. Use of innovative biomimetic scaffold in the treatment for large osteochondral lesions of the knee. Knee Surg Sports Traumatol Arthrosc [Internet]. 22 październik 2013 [cytowane 27 maj 2024]; Dostępne na: <http://link.springer.com/10.1007/s00167-013-2717-3>

36. Verdonk P, Dhollander A, Almqvist KF, Verdonk R, Victor J. Treatment of osteochondral lesions in the knee using a cell-free scaffold. Bone Jt J. marzec 2015;97-B(3):318–23.

37. Berruto M, Delcogliano M, De Caro F, Carimati G, Uboldi F, Ferrua P, i in. Treatment of Large Knee Osteochondral Lesions With a Biomimetic Scaffold: Results of a Multicenter Study of 49 Patients at 2-Year Follow-up. Am J Sports Med. lipiec 2014;42(7):1607–17.

38. Christensen BB, Foldager CB, Jensen J, Jensen NC, Lind M. Poor osteochondral repair by a biomimetic collagen scaffold: 1- to 3-year clinical and radiological follow-up. Knee Surg Sports Traumatol Arthrosc. lipiec 2016;24(7):2380–7.

39. Brix M, Kaipel M, Kellner R, Schreiner M, Apprich S, Boszotta H, i in. Successful osteoconduction but limited cartilage tissue quality following osteochondral repair by a cell-free multilayered nano-composite scaffold at the knee. Int Orthop. marzec 2016;40(3):625–32.

40. De Windt TS, Welsch GH, Brittberg M, Vonk LA, Marlovits S, Trattnig S, i in. Is Magnetic Resonance Imaging Reliable in Predicting Clinical Outcome After Articular Cartilage Repair of the Knee?: A Systematic Review and Meta-analysis. Am J Sports Med. lipiec 2013;41(7):1695–702.

41. Filardo G, Drobnic M, Perdisa F, Kon E, Hribernik M, Marcacci M. Fibrin glue improves osteochondral scaffold fixation: study on the human cadaveric knee exposed to continuous passive motion. Osteoarthritis Cartilage. kwiecień 2014;22(4):557–65.



REFERENCES