

Early Radiologic and Clinical Outcomes After Arthroscopic Rotator Cuff Repair Augmented with Demineralized Bone Fiber Implant

Aleksandar Banic¹, Kristine Italia^{1,2}, Christopher Child^{1,2}, Jashint Maharaj^{1,2},
Sarah Whithouse¹, **Ashish Gupta**^{1,2,3}

¹Queensland Unit for Advanced Shoulder Research (QUASR), ARC ITTC-JB, Queensland University of Technology, Brisbane, Australia

²Australian Shoulder Research Institute, Brisbane, Australia

³Greenslopes Private Hospital, Brisbane, Australia



Disclosures

QUASR- Funding from Australian Research Council, QUT, Stryker, Zimmer Biomet, Australian Biotechnologies, Materialise, Akunah


Australian Shoulder Research Fellowship (ASRI) - Funding from Stryker, Zimmer Biomet, Device Technology, Arthrex

AG: Consultant for Zimmer BIOMET, Device Tech, Sironix; Founder and CEO Akunah; hold shares with Tetrous



Introduction

- Healing at the enthesis after rotator cuff repair: weak link
- Re-tear rate after repair: can be as high as 94% in massive cuff tears
- **Aim:** To present the early radiological and clinical outcomes after arthroscopic repair of rotator cuff tears augmented with a novel demineralized bone fiber (DBF) implant (Enfix RC™, Tetra Tech Inc, Los Angeles, CA) at the footprint



Methods

- Patients who underwent arthroscopic rotator cuff repair with DBF augmentation
- **Comparison group:** patients who did not receive augmentation (matched according to age, sex, tendons involved, degree of retraction, follow-up period)
- **Outcomes**
 - Clinical Outcomes (preop, 6 months, 12 months)
 - Pain
 - Active range of motion
 - Isometric strength of rotator cuff measured by dynamometer
 - PROMs (ASES, Constant score, UCLA)
 - Radiologic outcomes (MRI at 6 months postop)
 - Tendon thickness at footprint
 - Sugaya classification
 - Healing

Results

- 31 patients in each group
- **Mean age: 54.8 years**
- **Mean follow-up: 8 months**
- Majority of patients with **at least 2-tendon tears with Patte 2-3 retraction and low-grade fatty infiltration** (comparable distribution between the 2 groups)



Table 1. Preoperative Patient and Rotator Cuff Tear Characteristics

Variables	Augmented Group	Control Group	p-value
n	31	31	
Age at surgery, y Mean (SD, range)	54.8 (6.1, 40-65)	54.8 (6.2, 42-64)	p=1.000
Sex (Male : Female)	24 (77.4%) : 7 (22.6%)	23 (74.2%) : 8 (25.8%)	p=0.767
Follow-up time, months Mean (SD, range)	7.8 (3.4, 6-14)	8.4 (3.4, 6-18)	p=0.388
Affected tendons based on Collin Classification			
Type A	1 (3.2%)	1 (3.2%)	p=0.617
Type B	2 (6.5%)	1 (3.2%)	
Type C	6 (19.4%)	4 (12.9%)	
Type D	15 (48.4%)	12 (38.7)	
Type E	2 (6.5%)	6 (19.4%)	
Type B+D [§]	3 (9.7%)	2 (6.5%)	
Type C+E [§]	0 (0%)	1 (3.2%)	
Type B+E [§]	0 (0%)	2 (6.5%)	p=0.478
Isolated supraspinatus	2 (6.5%)	2 (6.5%)	
Fatty Degeneration*			
SSC			
Goutallier 0-2	10/12 (83.3%)	11/11 (100%)	
Goutallier 3-4	2/12 (16.7%)	0/11 (0%)	
SSP			p=1.000
Goutallier 0-2	27/31 (87.1%)	27/31 (87.1%)	
Goutallier 3-4	4/31 (12.9%)	4/31 (12.9%)	p=1.000
ISP			
Goutallier 0-2	24/26 (92.3%)	25/27 (92.6%)	N/A
Goutallier 3-4	2/26 (7.7%)	2/27 (7.4%)	
TM			p=0.662
Goutallier 0-2	2/2 (100%)	9/9 (100%)	
Goutallier 3-4	0/2 (0%)	0/9 (0%)	
Tendon Retraction			
Patte 1	13 (41.9%)	10 (32.3%)	
Patte 2	4 (12.9%)	6 (19.4%)	
Patte 3	14 (45.2%)	15 (48.4%)	

SD, standard deviation; SSC, subscapularis; SSP, supraspinatus; ISP, infraspinatus; TM, teres minor.

[§]Not in the original Collin classification but the types have been combined to depict involved tendons.

*Only includes the ones with tears

Results


VAS

- Preop and postop pain scores were comparable between the 2 groups

Satisfaction rate

- Comparable between the 2 groups
- 96.8% in augmented group, 90.3% in control group, $p=0.612$)


*mean f/u = 8 months



Results

- Both groups improved from preop to postop
- **Augmented group had significantly higher Constant, ASES, and UCLA scores postoperatively**

*mean f/u = 8 months




Results

Active ROM

- Both groups had improvement in ROM in all planes
- Preop and postop ROM were comparable in both groups


*mean f/u = 8 months



Results

Absolute Strength

- Both groups had improvement strength
- Comparable preop and postop strengths




*mean f/u = 8 months

Results

Relative Strength

- Both groups achieved at least 90% strength of contralateral shoulder (except for supra strength in control group)
- Greater improvement in relative strength in augmented group but this was not statistically significant




*mean f/u = 8 months


Results

	With	Without	p value
Mean tendon thickness	5.3 mm	4.9mm	p=0.464
Healing	100%	100%	p=0.985
Tendon integrity			
<i>Sugaya 1</i>	3.7%	3.4%	
<i>Sugaya 2</i>	74.1%	72.4%	
<i>Sugaya 3</i>	22.2%	24.1%	
<i>Sugaya 4</i>	0%	0%	
<i>Sugaya 5</i>	0%	0%	

With augmentation



Without augmentation




*MRI at 6 months

Discussion

Demineralized Bone Fiber Implant

- Utilizes the biologic potential of DBM but in a form that has better handling characteristics and osteoconductivity, providing ease of use during arthroscopic rotator cuff repairs
- promotes healing and integration at the enthesis following rotator cuff repair through both osteoconductive and osteoinductive properties, and leads to development of a more functional four-zone enthesis with better histological and biomechanical properties



Conclusion

- Although both cohorts demonstrated comparable postoperative outcomes across most variables, the augmented group showed superior postoperative Constant, ASES, and UCLA scores and a trend toward rotator cuff strengths more closely resembling that of the non-operated shoulder at mean follow-up of 9 months
- The findings of this study demonstrates promising early results.
- However, extended follow-up and larger patient cohort is warranted to confirm these short-term outcomes presented in this study.